

MUPPADAI TRAINING ACADEMY – VELLORE

PREVIOUS YEAR QUESTIONS - 2017

NDA – MATHS

1. Let S be the set of all persons living in Delhi. We say that x, y in S are related if they were born in Delhi on the same day. Which one of the following is correct?
- A) The relation is an equivalent relation.
B) The relation is not reflexive but it is symmetric and transitive.
C) The relation is not symmetric but it is reflexive and transitive.
D) The relation is not transitive but it is reflexive and symmetric.
2. Let $A = \{1,2,3,4,5,6,7,8,9,10\}$. Then the number of subsets of A containing two or three elements is
- A) 45
B) 120
C) 165
D) 330
3. The value of $i^{2n} + i^{2n+1} + i^{2n+2} + i^{2n+3}$. Where $i = \sqrt{-1}$, is
- A) 0
B) 1
C) i
D) -i
4. If the difference between the roots of the equation $x^2 + kx + 1 = 0$ is strictly less than $\sqrt{5}$, Where $|k| \geq 2$, then k can be any element of the interval.
- A) $(-3, -2) \cup (2,3)$
B) $(-3,3)$
C) $(-3,-2) \cup (2,3)$
D) None of the above
5. If the roots of the equation $x^2 + px + q = 0$ are in the same ratio as those of the equation $x^2 + lx + m = 0$, then which one of the following is correct?
- A) $P^2m = l^2q$
B) $m^2P = l^2q$
C) $m^2P = q^2l$
D) $m^2P^2 = l^2q$
6. The value of $\left(\frac{1+i\sqrt{3}}{2}\right)^n + \left(\frac{1-i\sqrt{3}}{2}\right)^n$ Where n is not a multiple of 3 and $i = \sqrt{-1}$, is
- A) 1
B) -1
C) i
D) -i
7. Three – digit numbers are formed from the digits 1,2 and 3 in such a way that the digits are not repeated. What is the sum of such three digit numbers?
- A) 1233
B) 1322
C) 1323
D) 1332

MUPPADAI TRAINING ACADEMY – VELLORE

8. What is the sum of the series $0.3+0.33+0.333+\dots$ n terms?

- A) $\frac{1}{3}\left[n - \frac{1}{9}\left(1 - \frac{1}{10^n}\right)\right]$ B) $\frac{1}{3}\left[n - \frac{2}{9}\left(1 - \frac{1}{10^n}\right)\right]$
C) $\frac{1}{3}\left[n - \frac{1}{3}\left(1 - \frac{1}{10^n}\right)\right]$ D) $\frac{1}{3}\left[n - \frac{1}{9}\left(1 + \frac{1}{10^n}\right)\right]$

9. If $1, \omega, \omega^2$ are the cube roots of unity, then $(1 + \omega)(1 + \omega^2)(1 + \omega^3)(1 + \omega + \omega^2)$ is equal to

- A) -2 B) -1
C) 0 D) 2

10. If the sum of m terms of an AP is n and the sum of n terms is m, then the sum of (m+n) terms is

- A) mn B) m + n
C) $2(m+n)$ D) $-(m+n)$

11. The modulus and principal argument of the complex number $\frac{1+2i}{1-(1-i)^2}$ are respectively

- A) 1,0 B) 1,1
C) 2,0 D) 2,1

12. If the graph of a quadratic polynomial lies entirely above x-axis, then which one of the following is correct?

- A) Both the roots are real B) One root is real and the other is complex
C) Both the roots are complex D) Cannot say

13. If $|z + 4| \leq 3$, then the maximum value of $|z + 1|$ is

- A) 0 B) 4
C) 6 D) 10

14. The number of roots of the equation $z^2 = 2\bar{z}$ is

- A) 2 B) 3
C) 4 D) Zero

15. If $\cot \alpha$ and $\cot \beta$ are the roots of the equation $x^2 + bx + c = 0$ with $b \neq 0$, then the value of $\cot(\alpha + \beta)$ is

- A) $\frac{c-1}{b}$ B) $\frac{1-c}{b}$
C) $\frac{b}{c-1}$ D) $\frac{b}{1-c}$

16. The sum of the roots of the equation $x^2 + bx + c = 0$ (Where b and C are non-zero) is equal to the sum of the reciprocals of their squares. Then $\frac{1}{c}, b, \frac{c}{b}$ are in

- A) AP B) GP
C) HP D) None of the above

MUPPADAI TRAINING ACADEMY – VELLORE

17. The sum of the roots of the equation $ax^2+bx+C=0$ (Where a and C are non- zero) is equal to the sum of the reciprocals of their squares. Then a, ca^2 , c^2 are in
A) AP
B) GP
C) HP
D) None of the above
18. The value of $[C(7,0) + C(7,1)] + [C(7,1) + C(7,2)] + \dots + [C(7,6) + C(7,7)]$ is.
A) 254
B) 255
C) 256
D) 257
19. The number of different words (eight letter words) ending and beginning with a consonant which can be made out of the letters of the word 'EQUATION' is
A) 5200
B) 4320
C) 3000
D) 2160
20. The fifth term of an AP of n terms, whose sum is $n^2 - 2n$, is
A) 5
B) 7
C) 8
D) 15
21. The sum of all the two digit odd numbers is
A) 2475
B) 2530
C) 4905
D) 5049
22. The sum of the first n terms of the series $\frac{1}{2} + \frac{3}{4} + \frac{7}{8} + \frac{15}{16} + \dots$ is equal to
A) $2^n - n - 1$
B) $1 - 2^{-n}$
C) $2^n + n - 1$
D) $2^n - 1$
23. Consider the following in respect of sets A and B:
1. $(A - B) \cup B = A$ 2) $(A-B) \cup A = A$ 3) $(A - B) \cap B = \emptyset$ 4) $A \cup B = B$
Which of the above are correct?
A) 1,2 and 3
B) 2,3 and 4
C) 1,3 and 4
D) 1,2 and 4
24. In the binary equation $(1P101)_2 + (10q1)_2 = (100r00)_2$ Where p,q and r are binary digits, what are the possible values of p ,q and r respectively?
A) 0 , 1 , 0
B) 1, 1 ,0
C) 0, 0 , 1
D) 1, 0 , 1
25. If $S = \{x : x^2+1 = 0, x \text{ is real}\}$, then S is
A) $\{-1\}$
B) $\{0\}$
C) $\{1\}$
D) An empty set

MUPPADAI TRAINING ACADEMY – VELLORE

26. The expansion of $(x-y)^n$, $n \geq 5$ is done in the descending powers of x . If the sum of the fifth and sixth terms is zero, then $\frac{x}{y}$ is equal to

A) $\frac{n-5}{6}$
C) $\frac{5}{n-4}$

B) $\frac{n-4}{5}$
D) $\frac{6}{n-5}$

27. If $A = \begin{bmatrix} \alpha & 2 \\ 2 & \alpha \end{bmatrix}$ and $\det(A^3) = 125$, then α is equal to

A) ± 1
C) ± 3

B) ± 2
D) ± 5

28. If B is a non – singular matrix and A is a square matrix, then the value of $\det(B^{-1}AB)$ is equal to

A) $\det(B)$
C) $\det(B^{-1})$

B) $\det(A)$
D) $\det(A^{-1})$

29. If $a \neq b \neq c$, then one value of x which satisfies the equation

$$\begin{vmatrix} 0 & x-a & x-b \\ x+a & 0 & x-c \\ x+b & x+c & 0 \end{vmatrix} = 0$$

Is given by

A) a
C) c

B) b
D) 0

30. If $A = \begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix}$ then what is AA^T equal to (where A^T is the transpose of A) ?

A) Null matrix
C) A

B) Identity matrix
D) $-A$

31. The equations $x + 2y + 3z = 1$; $2x + y + 3z = 2$; $5x + 5y + 9z = 4$

A) Have the unique solution
C) are inconsistent

B) Have infinitely many solutions
D) None of the above

32. $A = \begin{bmatrix} x+y & y \\ x & x-y \end{bmatrix}$, $B = \begin{bmatrix} 3 \\ -2 \end{bmatrix}$ and $C = \begin{bmatrix} 4 \\ -2 \end{bmatrix}$ If $AB = C$, then What is A^2 equal to ?

A) $\begin{bmatrix} 4 & 8 \\ -4 & -16 \end{bmatrix}$
C) $\begin{bmatrix} -4 & -8 \\ 4 & 12 \end{bmatrix}$

B) $\begin{bmatrix} 4 & -4 \\ 8 & -16 \end{bmatrix}$
D) $\begin{bmatrix} -4 & -8 \\ 8 & 12 \end{bmatrix}$

MUPPADAI TRAINING ACADEMY – VELLORE

33. What is the value of the determinant $\begin{vmatrix} 1 & 1 & 1 \\ 1 & 1+xyz & 1 \\ 1 & 1 & 1+xyz \end{vmatrix}$?

- A) $1+x+y+z$
 C) $x^2y^2z^2$

- B) $2xyz$
 D) $2x^2y^2z^2$

34. If $\begin{vmatrix} x & y & 0 \\ 0 & x & y \\ y & 0 & x \end{vmatrix} = 0$, then which one of the following is correct?

- A) $\frac{x}{y}$ Is one of the cube roots of unity.
 B) x is one of the cube roots of unity
 C) Y is one of the cube roots of unity
 D) $\frac{x}{y}$ is one of the cube roots of -1.

35. Consider the set A of all matrices order 3×3 with entries 0 or 1 only. Let B be the subset of A consisting of all matrices whose determinant is 1. Let C be the subset of A consisting of all matrices whose determinant is -1. Then which one of the following is correct?

- A) C is empty
 B) B has as many elements as C
 C) $A = B \cup C$
 D) B has thrice as many elements as C.

36. If $A = \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{bmatrix}$, then what is A^3 equal to ?

- A) $\begin{bmatrix} \cos 3\theta & \sin 3\theta \\ -\sin 3\theta & \cos 3\theta \end{bmatrix}$
 C) $\begin{bmatrix} \cos 3\theta & -\sin 3\theta \\ \sin 3\theta & \cos 3\theta \end{bmatrix}$

- B) $\begin{bmatrix} \cos^3\theta & \sin^3\theta \\ -\sin^3\theta & \cos^3\theta \end{bmatrix}$
 D) $\begin{bmatrix} \cos^3\theta & -\sin^3\theta \\ \sin^3\theta & \cos^3\theta \end{bmatrix}$

37. What is the order of $\begin{bmatrix} a & h & g \\ h & b & f \\ g & f & c \end{bmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix}$?

- A) 3×1
 B) 1×1
 C) 1×3
 D) 3×3

38. If $a = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, then the value of A^4 is

- A) $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$
 B) $\begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}$
 C) $\begin{pmatrix} 0 & 0 \\ 1 & 1 \end{pmatrix}$
 D) $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$

MUPPADAI TRAINING ACADEMY – VELLORE

39. If $\sin A = \frac{3}{5}$, where $450^\circ < A < 540^\circ$, then $\cos \frac{A}{2}$ is equal to

A) $\frac{1}{\sqrt{10}}$

B) $-\sqrt{\frac{3}{10}}$

C) $\frac{\sqrt{3}}{\sqrt{10}}$

D) None of the above

40. What is $\frac{1}{\sin 10^\circ} - \frac{\sqrt{3}}{\cos 10^\circ}$ equal to ?

A) 0

B) 1

C) 2

D) 4

41. From the top of a lighthouse, 100 m high, the angle of depression of a boat is $\tan^{-1}\left(\frac{5}{12}\right)$. What is the distance between the boat and the lighthouse?

A) 120 m

B) 180m

C) 240m

D) 360m

42. The maximum value of $\sin \left[x + \frac{\pi}{6} \right] + \cos \left[x + \frac{\pi}{6} \right]$ in the interval $\left[0, \frac{\pi}{2} \right]$ is attained at

A) $\frac{\pi}{12}$

B) $\frac{\pi}{6}$

C) $\frac{\pi}{3}$

D) $\frac{\pi}{2}$

43. If $K = \sin \left[\frac{\pi}{18} \right] \sin \left[\frac{5\pi}{18} \right] \sin \left[\frac{7\pi}{18} \right]$, then What is the value of K?

A) $\frac{1}{2}$

B) $\frac{1}{4}$

C) $\frac{1}{8}$

D) $\frac{1}{16}$

44. The expression $\frac{\sin \alpha + \sin \beta}{\cos \alpha + \cos \beta}$ is equal to

A) $\tan \left[\frac{\alpha + \beta}{2} \right]$

B) $\cot \left[\frac{\alpha + \beta}{2} \right]$

C) $\sin \left[\frac{\alpha + \beta}{2} \right]$

D) $\cos \left[\frac{\alpha + \beta}{2} \right]$

45. If $\sin \theta = 3 \sin(\theta + 2\alpha)$, then the value of $\tan(\theta + \alpha) + 2 \tan \alpha$ is equal to

A) -1

B) 0

C) 1

D) 2

46. What is the value of $\tan 18^\circ$?

A) $\frac{\sqrt{5}-1}{\sqrt{10+2\sqrt{5}}}$

B) $\frac{\sqrt{5}-1}{\sqrt{10+\sqrt{5}}}$

C) $\frac{\sqrt{10+2\sqrt{5}}}{\sqrt{5}-1}$

D) $\frac{\sqrt{10+\sqrt{5}}}{\sqrt{5}-1}$

MUPPADAI TRAINING ACADEMY – VELLORE

54. The two circles $x^2 + y^2 = r^2$ and $x^2 - y^2 - 10x + 16 = 0$ intersect at two distinct points. Then which one of the following is correct?
A) $2 < r < 8$
B) $r = 2$ or $r = 8$
C) $r < 2$
D) $r > 2$
55. What is the equation of the circle which passes through the points (3,-2) and (-2,0) and having its centre on the line $2x - y - 3 = 0$?
A) $x^2 + y^2 + 3x + 2 = 0$
B) $x^2 + y^2 + 3x + 12y + 2 = 0$
C) $x^2 + y^2 + 2x = 0$
D) $x^2 + y^2 = 5$
56. What is the ratio in which the point C $[-\frac{2}{7}, -\frac{20}{7}]$ divides the line joining the points A(-2,-2) and B(2,-4)?
A) 1:3
B) 3:4
C) 1:2
D) 2:3
57. What is the equation of the ellipse having foci $(\pm 2, 0)$ and the eccentricity $\frac{1}{4}$?
A) $\frac{x^2}{64} + \frac{y^2}{60} = 1$
B) $\frac{x^2}{60} + \frac{y^2}{64} = 1$
C) $\frac{x^2}{20} + \frac{y^2}{24} = 1$
D) $\frac{x^2}{24} + \frac{y^2}{20} = 1$
58. What is the equation of the straight line parallel to $2x + 3y + 1 = 0$ and passes through the point (-1,2)?
A) $2x + 3y - 4 = 0$
B) $2x + 3y - 5 = 0$
C) $x + y - 1 = 0$
D) $3x - 2y + 7 = 0$
59. What is the acute angle between the pair of straight lines $\sqrt{2}x + \sqrt{3}y = 1$ and $\sqrt{3}x + \sqrt{2}y = 2$?
A) $\tan^{-1}(\frac{1}{2\sqrt{6}})$
B) $\tan^{-1}(\frac{1}{\sqrt{2}})$
C) $\tan^{-1}(3)$
D) $\tan^{-1}(\frac{1}{\sqrt{3}})$
60. If the centroid of a triangle formed by (7,x), (y,-6) and (9,10) is (6,3), then the values of x and y are respectively.
A) 5,2
B) 2,5
C) 1,0
D) 0,0
61. A straight line with direction cosines (0,1,0) is
A) Parallel to x-axis
B) Parallel to y-axis
C) Parallel to z axis
D) Equally inclined to all the axes

MUPPADAI TRAINING ACADEMY – VELLORE

62. $(0,0,0), (\alpha,0,0), (0,b,0)$ and $(0,0,C)$ are four distinct points. What are the coordinates of the point which is equidistant from the four points?

- A) $[\frac{a+b+c}{3}, \frac{a+b+c}{3}, \frac{a+b+c}{3}]$ B) (a,b,c)
C) $(\frac{a}{2}, \frac{b}{2}, \frac{c}{2})$ D) $[\frac{a}{3}, \frac{b}{3}, \frac{c}{3}]$

63. The points $P(3,2,4), Q(4,5,2), R(5,8,0)$ and $S(2,-1,6)$ are

- A) Vertices of a rhombus which is not a square
B) Non- Coplanar
C) Collinear
D) Coplanar but not collinear.

64. The line passing through the points $(1,2,-1)$ and $(3,-1,2)$ meets the yz -plane at which one of the following points?

- A) $[0, \frac{7}{2}, \frac{5}{2}]$ B) $[0, \frac{7}{2}, \frac{1}{2}]$
C) $[0, -\frac{7}{2}, -\frac{5}{2}]$ D) $[0, \frac{7}{2}, -\frac{5}{2}]$

65. Under which one of the following conditions are the lines $x = ay + b; z = cy + d$ and $x = ey + f; z = gy + h$ perpendicular?

- A) $ae + cg - 1 = 0$ B) $ae + bf - 1 = 0$
C) $ae + cg + 1 = 0$ D) $ag + ce + 1 = 0$

66. If $\vec{a} = i - j + k, \vec{b} = 2i + 3j + 2k$ and $\vec{c} = i + mj + nk$ are three coplanar vectors and $|\vec{c}| = \sqrt{6}$, then which one of the following is correct?

- A) $m = 2$ and $n = \pm 1$ B) $m = \pm 2$ and $n = -1$
C) $m = 2$ and $n = -1$ D) $m = \pm 2$ and $n = 1$

67. Let ABCD be a parallelogram whose diagonals intersect at P and let O be the origin. What is $\vec{OA} + \vec{OB} + \vec{OC} + \vec{OD}$ equal to?

- A) $2\vec{OP}$ B) $4\vec{OP}$
C) $6\vec{OP}$ D) $8\vec{OP}$

68. ABCD is a quadrilateral whose diagonals are AC and BD which one of the following is correct?

- A) $\vec{BA} + \vec{CD} = \vec{AC} + \vec{DB}$ B) $\vec{BA} + \vec{CD} - \vec{BD} + \vec{CA}$
C) $\vec{BA} + \vec{CD} = \vec{AC} + \vec{BD}$ D) $\vec{BA} + \vec{CD} = \vec{BC} + \vec{AD}$

MUPPADAI TRAINING ACADEMY – VELLORE

69. If $\vec{a} \times \vec{b} = \vec{c}$ and $\vec{b} \times \vec{c} = \vec{a}$, then which one of the following is correct?

- A) $\vec{a}, \vec{b}, \vec{c}$ are orthogonal in pairs and $|\vec{a}| = |\vec{c}|$ and $|\vec{b}| = 1$
- B) $\vec{a}, \vec{b}, \vec{c}$ are non – orthogonal to each other
- C) $\vec{a}, \vec{b}, \vec{c}$ are orthogonal in pairs but $|\vec{a}| \neq |\vec{c}|$
- D) $\vec{a}, \vec{b}, \vec{c}$ are orthogonal in pairs but $|\vec{b}| \neq 1$

70. If $\vec{a} = 2\hat{i} + 3\hat{j} + 4\hat{k}$, $\vec{b} = 3\hat{i} + 2\hat{j} - \tau\hat{k}$ are perpendicular, then what is the value of τ ?

- A) 2
- B) 3
- C) 4
- D) 5

71. What is $\lim_{x \rightarrow 0} \frac{e^x - (1+x)}{x^2}$ equal to ?

- A) 0
- B) $\frac{1}{2}$
- C) 1
- D) 2

72. What is $\int_0^{\frac{\pi}{2}} \frac{d\theta}{1+\cos\theta}$ equal to?

- A) $\frac{1}{2}$
- B) 1
- C) $\sqrt{3}$
- D) None of the above

73. What is $\int \frac{dx}{x(x^7+1)}$ equal to ?

- A) $\frac{1}{2} \ln \left| \frac{x^7-1}{x^7+1} \right| + c$
- B) $\frac{1}{7} \ln \left| \frac{x^7+1}{x^7} \right| + c$
- C) $\ln \left| \frac{x^7-1}{7x} \right| + c$
- D) $\frac{1}{7} \ln \left| \frac{x^7}{x^7+1} \right| + c$

74. The function $f: X \rightarrow y$ defined by $f(x) = \cos x$, where $x \in X$, is one – one and onto if x and y are respectively equal to

- A) $[0, \pi]$ and $[-1, 1]$
- B) $[-\frac{\pi}{2}, \frac{\pi}{2}]$ and $[-1, 1]$
- C) $[0, \pi]$ and $[-1, 1]$
- D) $|0, x|$ and $|0, 1|$

75. If $f(x) = \frac{x}{x-1}$ then what is $\frac{f(a)}{f(a+1)}$ equal to?

- A) $f - \left(\frac{a}{a+1}\right)$
- B) $f(a^2)$
- C) $f\left(\frac{1}{a}\right)$
- D) $f|-a|$

MUPPADAI TRAINING ACADEMY – VELLORE

76. What is $\int \frac{(x^{x-1} + e^{x-1})dx}{x^e + e^x}$ equal to?

A) $\frac{x^2}{2} + c$

B) $\ln(x + e) + c$

C) $\ln(x^e + e^x) + c$

D) $\frac{1}{e} \ln(x^e + e^x) + e$

77. Let $f: |-6, 6| \rightarrow R$ be defined by $f(x) = x^2 - 3$, Consider the following:

1. $(f \circ f \circ f)(-1) = (f \circ f \circ f)(1)$

2. $(f \circ f \circ f)(-1)(-4)(f \circ f \circ f)(1)$

Which of the above is/are correct?

A) 1 only

B) 2 only

C) Both 1 and 2

D) Neither 1 nor 2

78. Let $f(x) = px + q$ and $g(x) = mx + n$. Then $f(g(x)) = g(f(x))$ is equivalent to

A) $f(p) = g(m)$

B) $f(g) = g(n)$

C) $f(n) = g(q)$

D) $f(m) = g(p)$

79. If $f(x) = \sqrt{9 - x^2}$, then what is $\lim_{x \rightarrow 1} \frac{f(x) - f(1)}{x - 1}$ equal to?

A) $-\frac{1}{4\sqrt{2}}$

B) $\frac{1}{8}$

C) $-\frac{1}{2\sqrt{2}}$

D) $\frac{1}{2\sqrt{2}}$

80. What is $\frac{d^2x}{dy^2}$ equal to?

A) $-\left(\frac{d^2y}{dx^2}\right)^{-1} \left(\frac{dy}{dx}\right)^{-3}$

B) $\left(\frac{d^2y}{dx^2}\right)^{-1} \left(\frac{dy}{dx}\right)^{-3}$

C) $-\left(\frac{d^2y}{dx^2}\right) \left(\frac{dy}{dx}\right)^{-3}$

D) $\left(\frac{d^2y}{dx^2}\right)^{-1}$

81. Let

$$f(x): \begin{cases} x, x \text{ is rational} \\ 0, x \text{ is irrational} \end{cases}$$

and

$$g(x): \begin{cases} 0, x \text{ is rational} \\ x, x \text{ is irrational} \end{cases}$$

If $f: R \rightarrow R$ and $g: R \rightarrow R$, then $(f - g)$ is

A) one-to-one and into

B) neither one-to-one nor onto

C) many-to-one and onto

D) one-to-one and onto

MUPPADAI TRAINING ACADEMY – VELLORE

82. What is the length of the longest interval in which the function $f(x) = 3 \sin x - 4 \sin^2 x$ is increasing?

- A) $\frac{\pi}{3}$ B) $\frac{\pi}{2}$
C) $\frac{3\pi}{2}$ D) x

83. If $xdy = y(dx + ydy)$; $y(1) = 1$ and $y(x) > 0$, then what is $y(-3)$ equal to?

- A) 3 B) 2
C) 1 D) 0

84. What is maximum value of the function $f(x) = 4 \sin^2 x + 1$?

- A) 5 B) 3
C) 2 D) 1

85. Let $f(x)$ be an indefinite integral of $\sin^2 x$. Consider the following statements:

Statement : 1

The function $f(x)$ satisfies $f(x + \pi) - f(x)$ for all real x

Statement: 2

$\sin^2(x - 1) = \sin^2 x$ for all real x .

Which one of the following is correct in respect of the above statements?

- A) Both the statements are true and Statement 2 is the correct explanation of Statement 1
B) Both the Statements are true but Statement 2 is not the correct explanation of Statement 1
C) Statement 1 is true but Statement 2 is false
D) Statement 1 is false but Statement 2 is true

86. What are the degree and order respectively of the differential equation?

$$y = x \left(\frac{dy}{dx} \right)^3 + \left(\frac{dx}{dy} \right)^2 ?$$

- A) 1, 2 B) 2, 1
C) 1, 4 D) 4, 1

87. What is the differential equation corresponding to $y^2 - 2ay + x^2 = a^2$ by eliminating a ?

- A) $(x^2 - 2y^2)p^2 - 4pxy + x^2 = 0$
B) $(x^2 - 2y^2)p^2 + 4pxy - x^2 = 0$
C) $(x^2 + 2y^2)p^2 - 4pxy - x^2 = 0$
D) $(x^2 + 2y^2)p^2 - 4pxy + x^2 = 0$

Where $p = \frac{dy}{dx}$

MUPPADAI TRAINING ACADEMY – VELLORE

88. What is the general solution of the differential equation

$$ydx - (x + 2y^2)dy = 0?$$

A) $x = y^2 + cy$

B) $x = 2xy^2$

C) $x - 2y^2 + cy$

D) None of the above

89. Let

$$f(x + y) =$$

$f(x)f(y)$ for all x and y . Then what is $f'(5)$ equal to where $f'(x)$ is the derivation of $f(x)$?

A) $f(5)f'(0)$

B) $f(5)-f'(0)$

C) $f(5)f(0)$

D) $f(5)+f'(0)$

90. If $f(x)$ and $g(x)$ are continuous function satisfying $f(x) = f(a-x)$ and $g(x)+f(a-x)=2$, then what is

A) $\int_0^a g(x)dx$

B) $\int_0^a f(x)dx$

C) $2\int_0^a f(x)dx$

D) 0

91. What is the solution of the differential equation in $\left(\frac{dy}{dx}\right) - a = 0$?

A) $y = xe^a + c$

B) $x = ye^a + c$

C) $y = 1nx + c$

D) $x=1ny+c$

92. Let $f(x)$ be defined as follows: -

$$f(x) = \begin{cases} 2x+1, & -3 < x < -2 \\ x-1, & -2 \leq x < 0 \\ x+2, & 0 \leq x < 1 \end{cases}$$

Which one of the following statements is correct in respect of the above function?

A) It is discontinuous at $x = -2$ but continuous at every other point.

B) It is continuous only in the interval $(-3, -2)$

C) It is discontinuous at $x=0$ but continuous at every other point.

D) It is discontinuous at every point.

93. Consider the following statements:-

1. If $\lim_{x \rightarrow a} f(x)$ and $\lim_{x \rightarrow a} g(x)$ both exist, then $\lim_{x \rightarrow a} (f(x)g(x))$ exists.

2. If $\lim_{x \rightarrow a} (f(x)g(x))$ exists, then both $\lim_{x \rightarrow a} f(x)$ and $\lim_{x \rightarrow a} g(x)$ must exist.

Which of the above statements is/are correct?

A) 1 only

B) 2 only

C) Both 1 and 2

D) Neither 1 nor 2

MUPPADAI TRAINING ACADEMY – VELLORE

94. Which one of the following functions is neither even nor odd?

- A) $x^2 + 1$ B) $x + \frac{3}{x}$
C) (x) D) $x^2(x - 3)$

95. What is the derivative of $\log_{10}(5x^2 + 3)$ with respect to x ?

- A) $\frac{x \log_{10} e}{5x^2+3}$ B) $\frac{2x \log_{10} e}{5x^2+3}$
C) $\frac{10x \log_{10} e}{5x^2+3}$ D) $\frac{x \log_e 10}{5x^2+3}$

96. Let $f(a) = \frac{a-1}{a+1}$

Consider the following:-

1. $f(2a) = f(a) + 1$
2. $f\left(\frac{1}{a}\right) = -f(a)$

Which of the above is/are correct?

- A) 1 only B) 2 only
C) Both 1 and 2 D) Neither 1 nor 2

97. What is the maximum area of a triangle that can be inscribed in a circle of radius a ?

- A) $\frac{3a^2}{4}$ B) $\frac{a^2}{2}$
C) $\frac{3\sqrt{3}a^2}{4}$ D) $\frac{\sqrt{3}a^2}{4}$

98. Let $f(x) = x + \frac{1}{x}$ Where $x \in (0,1)$. Then which one of the following is correct?

- A) $f(x)$ fluctuates in the interval
B) $f(x)$ increases in the interval
C) $f(x)$ decreases in the interval
D) None of the above

99. Suppose the function

$f(x) = x^n, n \neq 0$ is differentiable for all x . Then n can be any element of the interval.

- A) $(1, \infty)$ B) $(0, \infty)$
C) $\left(\frac{1}{2}, \infty\right)$ D) None of the above

100. What is $\int_{-1}^{0^2} \left| \frac{1nx}{x} \right| dx$ equal to?

- A) $\frac{3}{2}$ B) $\frac{5}{2}$
C) 3 D) 4

MUPPADAI TRAINING ACADEMY – VELLORE

101. The variance of 20 observations is 5. If each observation is multiplied by 3, then what is the new variance of the resulting observations?
- A) 5
B) 10
C) 15
D) 45
102. The mean of a group of 100 observations was found to be 20. Later it was found that four observations were incorrect, which were recorded as 21, 21, 18 and 20, what is the mean if the incorrect observations are omitted?.
- A) 18
B) 20
C) 21
D) 22
103. A Committee of two persons is constituted from two men and two women. What is the probability that the committee will have only women?
- A) $\frac{1}{6}$
B) $\frac{1}{3}$
C) $\frac{1}{2}$
D) $\frac{2}{3}$
104. A question is given to three students A, B and C whose chances of solving it are $\frac{1}{2}$, $\frac{1}{3}$ and $\frac{1}{4}$ respectively. What is the probability that the question will be solved?
- A) $\frac{1}{24}$
B) $\frac{1}{4}$
C) $\frac{1}{2}$
D) $\frac{2}{3}$
105. The mean weight of 150 students in a certain class is 60kg. The mean weight of boys in the class is 70 kg and that of girls is 55kg. What is the number of boys in the class?
- A) 50
B) 55
C) 60
D) 100
106. For two dependent events A and B, it is given that $P(A) = 0.2$ and $P(B) = 0.5$. If $A \subseteq B$, then the values of conditional probabilities $P(A|B)$ and $P(B|A)$ are respectively
- A) $\frac{2}{5}, \frac{3}{5}$
B) $\frac{2}{5}, 1$
C) $1, \frac{2}{5}$
D) Information is insufficient
107. A point is chosen at random inside a circle. What is the probability that the point is closer to the centre of the circle than to its boundary?
- A) $\frac{1}{5}$
B) $\frac{1}{4}$
C) $\frac{1}{3}$
D) $\frac{1}{2}$

MUPPADAI TRAINING ACADEMY – VELLORE

114. For given statistical data, the graphs for less than o give and more than o give are drawn. If the point at which the two curves intersect is P, then abscissa of point P gives the value of which one of the following measures of central tendency?
- A) Median
B) Mean
C) Mode
D) Geometric mean
115. Consider the following statements:-
1. Two events are mutually exclusive if the occurrence of one event prevents the occurrence of the other.
 2. The probability of the union of two mutually exclusive events is the sum of their individual probabilities.
- Which of the above statements is/are correct?
- A) 1 only
B) 2 only
C) Both 1 and 2
D) Neither 1 nor 2
116. If the regression coefficient of x on y and y on x are $-1/2$ and $-1/8$ respectively, then what is the correlation coefficient between x and y?
- A) $-1/4$
B) $-1/16$
C) $1/16$
D) $1/4$
117. A sample of 5 observations has mean 32 and median 33. Later it is found that an observation was recorded incorrectly as 40 instead of 35. If we correct the data. Then which one of the following is correct?
- A) The mean and median remain the same
B) The median remains the same but the mean will decrease
C) The mean and median both will decrease.
D) The mean remains the same but median will decrease.
118. If two fair dice are thrown, then what is the probability that the sum is neither 8 nor 9?
- A) $1/6$
B) $1/4$
C) $3/4$
D) $5/6$
119. Let A and B are two mutually exclusive events with $P(A) = 1/3$ $P(B) = 1/4$ what is the value of $P(\bar{A} \cap \bar{B})$
- A) $1/6$
B) $1/4$
C) $1/3$
D) $5/12$
120. The mean and standard deviation of a binomial distribution are 12 and 2 respectively. What is the number of trials?
- A) 2
B) 12
C) 18
D) 24